

USE ONLY AS DIRECTED

Bravecto® Spot-On should be administered in accordance with the following table

Weight of dog/cat (kg)	Pipette size to be used	Volume (mℓ)	Fluralaner (mg)
2 - 4,5	Miniature dogs	0,4	112,5
> 4,5 - 10	Small dogs	0,89	250
> 10 - 20	Medium dogs	1,79	500
> 20 - 40	Large dogs	3,57	1 000
> 40 - 56	Extra-large dogs	5,0	1 400
> 56	The appropriate combination of pipettes should be used.		
1,2 - 2,8	Small cats	0,4	112,5
> 2,8 - 6,25	Medium cats	0,89	250
> 6,25 - 12,5	Large cats	1,79	500

Within each weight range a whole pipette must be used.


For cats weighing more than 12,5 kg, use a combination of 2 pipettes that most closely matches the body weight.

Method of administration

Step 1:

Immediately before use, open the sachet and remove the pipette. The pipette should be held by the upper part of the tail base (crimped end) or by the upper rigid portion below the cap in an upright position (tip up) before opening it. The cap should be rotated clockwise or

counter clockwise one full turn. **The cap will stay on the pipette; it should not be removed.** The pipette is open and ready for application when the breaking of the seal is felt.



Step 2:

The cat should be standing or lying with its back horizontally during application. Part the fur at the administration site. Place the pipette tip vertically against the skin between the shoulder blades of the dog or pipette tip on the base of the skull of the cat.

Step 3:

Squeeze the pipette gently and apply the entire contents directly to the cat's skin. The product should be applied in one spot (cats up to 6,25 kg body weight) or two spots (cats weighing more than 6,25 kg body weight). If two spots are needed, the first spot should be applied at the base of the skull and the second one between the shoulder blades.

It is important to deliver the solution onto an area that cannot be easily licked by the cat. As cats are fastidious groomers, they may inadvertently ingest product if easily reached. Ingestion may cause adverse reactions and will remove product from the coat. If multiple cats are treated and may mutually groom, it is recommended that cats are separated while product dries.