

Use only as directed.

Administer orally at monthly intervals to provide the recommended dosage of 2,5 mg/kg of afoxolaner and 0,50 mg/kg of milbemycin oxime per kg bodyweight.

For dogs above 60 kg, appropriate combinations of chewable tablets are used.

NEXGARD® SPECTRA is only recommended for use in dogs and puppies 8 weeks of age and older and dogs weighing more than 2 kg.

| PRESENTATION                | TABLET SIZE ( g) | AFOXOLANER (mg /TABLET) | MILBEMYCIN OXIME (mg /TABLET) |
|-----------------------------|------------------|-------------------------|-------------------------------|
| Very small dogs (2–3,5 kg)  | 0,5              | 9,38                    | 1,88                          |
| Small dogs (3,6-7,5 kg)     | 1                | 18,75                   | 3,75                          |
| Medlum dogs (7,6–15 kg)     | 2                | 37,5                    | 7,5                           |
| Large dogs (15–30 kg)       | 4                | 75,0                    | 15,0                          |
| Extra Large dogs (30-60 kg) | 8                | 150,0                   | 30,0                          |

The following dosing table serves as a guide:

#### METHOD OF ADMINISTRATION

For the treatment and prevention of tick and flea infestation in dogs:

Administer the chewable tablets appropriate for your dog's weight at monthly intervals. Because dogs find **NEXGARD® SPECTRA** palatable, the product can be offered to the dog by hand. Alternatively, chewable tablets can be added to a small amount of dog food, or dosed manually.