

Administer **REVOLUTION** in accordance with the following tables:

REVOLUTION is available in tube sizes for dogs and cats of different mass.

**For cats and kittens:**

<b>Mass (kg)</b>	<b>Package colour</b>	<b>Potency (mg/ ml )</b>	<b>Administered volume (nominal tube sizes - ml)</b>
Up to 2,5	Mauve	60	0,25
2,6-7,5	Blue	60	0,75
>7,5		60	Appropriate combination of tubes

**For dogs and puppies:**

<b>Mass (kg)</b>	<b>Package colour</b>	<b>Potency (mg/ ml)</b>	<b>Administered volume (nominal tube sizes - ml)</b>
Up to 2,5	Mauve	60	0,25
2,6-5,0	Purple	120	0,25
5,1-10,0	Brown	120	0,5
10,1-20,0	Red	120	1,0
20,1-40,0	Teal	120	2,0
>40		120	Appropriate combination of tubes

Depress the cap to puncture the seal on the REVOLUTION tube; then remove the cap to administer product. Administer the product directly to the skin without massaging into the site. Part the hair on the back of the animal at the base of the neck in front of the shoulder blades until the skin is visible. Place the tip of the tube on the skin and squeeze the tube to empty its entire contents directly onto the skin in one spot. Avoid contact between the product and fingers. Do not apply when the hair coat is wet. Bathing or shampooing the animal 2 or more hours after treatment will not reduce the efficacy of REVOLUTION.

**EFFICACY:**

Adult fleas on dogs and cats are killed for one month following administration of REVOLUTION according to the recommended dosing schedule. In addition, flea eggs exposed to this REVOLUTION dosing schedule are not viable and flea reproduction is stopped. By killing adult fleas, REVOLUTION controls signs of flea infestation including flea allergy dermatitis. Ear mites (*O. cynotis*) in dogs and cats, sarcoptic mange mites (*S. scabiei*) in dogs, and intestinal hookworms (*A. tubaeforme*) in cats and roundworms in cats and dogs are killed after a single dose of REVOLUTION and controlled by monthly dosing. REVOLUTION should be applied once as a single treatment for lice infestation. Monthly use of REVOLUTION will control any subsequent lice infestations. Monthly administration of REVOLUTION prevents heartworm disease by controlling the microfilaria of *D. immitis* in dogs and cats. Ticks are killed when engorging on dogs and cats that have been treated with REVOLUTION. In heavy tick infestations, complete efficacy may not be achieved after the first dose therefore REVOLUTION should be administered on a monthly basis with one additional dose administered two weeks after the first dose with monthly dosing continued thereafter.