

WARNINGS

- Keep out of reach of children, uninformed persons and animals.
- Mild and transient skin reactions at the application site (itching, hairloss) may occur on rare occasions. These signs disappear without further treatment.
- If the cat licks the application site after treatment, temporary excessive salivation can be observed.
- Oral ingestion of BROADLINE® may also result in transient neurological signs. Signs resolve without treatment within 24 hours. Correct application will minimise the occurrence of such events.
- A temporary clumping or spiking of the hair may be seen at the application site after treatment.
- Pregnancy and lactation: The safety of the veterinary medicinal product has not been established during pregnancy and lactation. Laboratory studies with the individual ingredients in rats and rabbits have not produced teratogenic, foetotoxic or maternotoxic effects. Use only according to the risk-benefit assessment by the attending veterinarian.
- Interaction with other medicinal products: No interactions between BROADLINE® and routinely used veterinary medicinal products, or medical or surgical procedures have been observed.

- During treatment with BROADLINE®, concurrent administration of other antiparasitic macrocyclic lactone should be based upon the risk/benefit assessment of the attending veterinarian, since the potential of interactions in such cases has not been investigated.
- Avoid contact with the cat's eyes.
- After treatment with BROADLINE®, ticks will generally be killed within 48 hours after infestation without having a blood meal. However the attachment of single ticks cannot be excluded and transmission of infectious diseases cannot thus be completely excluded.
- No data on the effect of bathing/shampooing on the efficacy of BROADLINE® on cats is available.
- However, brief contact of the animal with water on one or two occasions within the month following application is unlikely to significantly reduce the efficacy of BROADLINE®.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Special precautions for use in animals:
- In the absence of available data, do not repeat the treatment at intervals of less than 2 weeks; the treatment of kittens < 7 weeks and/or below 0.6 kg should be based on a risk-benefit assessment.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

- Do not drink, eat or smoke while handling BROADLINE®.
- Wash hands after use.
- Used applicators should be disposed of immediately. Stored applicators must be kept in the intact blister package.
- In case of accidental spillage onto skin, wash off immediately with soap and water. In case of accidental eye exposure, flush the eyes carefully with