

Warnings:

- For animal use
- Do not use in puppies younger than 6 weeks of age.
- Do not use **Milbemax™ Chewable Tablets** for Dogs more Than 5 kg in dogs with a body mass of less than 5 kg.
- The margin of safety of **Milbemax™ Chewable** in Collies or related dog breeds is less than other breeds and therefore the recommended dose should be strictly observed in these breeds.
- In heartworm-risk areas, or where dogs have been travelling to or from heartworm-risk regions, a veterinary consultation is recommended to exclude the presence of any concurrent heartworm (*Dirofilaria immitis*) infestation. In the case of a positive diagnosis, adulticidal therapy is indicated before administering **Milbemax™ Chewable**.
- *Echinococcus* represents a hazard for humans (zoonosis). In cases of echinococcosis, specific guidelines on the treatment, follow-up and safeguarding of such persons should be followed.
- In severely debilitated dogs or individuals with seriously compromised kidney or liver function, the use of **Milbemax™ Chewable** is recommended only according to a risk/benefit assessment by the responsible veterinarian as no studies have been performed in such cases.
- At 10 times and higher dose rates (overdosing), mild transient clinical signs of mydriasis, depression, drooling, paresis, trembling and/or a staggered gait may occur in some dogs. Such symptoms will subside spontaneously within a day. No antidote is known.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

Precautions:

- The concurrent use of **Milbemax™ Chewable** with the macrocyclic lactone selamectin is well tolerated. No interactions were observed when the recommended dose of selamectin was administered concomitantly with **Milbemax™ Chewable** at the recommended dose. In the absence of studies, caution should be exercised when other macrocyclic lactones are administered concomitantly with **Milbemax™ Chewable**.
- Treatment of dogs with a high number of circulating microfilariae can sometimes lead to the appearance of hypersensitivity reactions, such as pale mucous membranes, vomiting, trembling, laboured breathing or excessive salivation. These reactions are associated with the release of proteins from dead or dying microfilariae and are not a direct toxic effect of the product. The use in dogs suffering from microfilaraemia is therefore not recommended.
- Wash hands after use.
- Do not contaminate water courses, as milbemycin may be harmful to fish and other aquatic organisms.

First aid: in the event of accidental ingestion of tablets, particularly by a child, seek medical advice immediately and show this package insert to the physician.