

WARNINGS

- Keep out of reach of children, uninformed persons and animals.
- After treatment with NEXGARD®, most ticks will generally be killed within 48 hours after infestation. However the attachment of single ticks cannot be excluded and so transmission of infectious diseases cannot be completely excluded if conditions are unfavourable.
- NEXGARD® may cause vomiting and diarrhoea in some dogs.
- Use during pregnancy, lactation or breeding: No data have been generated in dogs used for breeding. However, in laboratory species (rats and rabbits), afoxolaner has not produced any evidence of teratogenic effects, nor any adverse effect on the reproductive capacity in males and females. Safety in pregnant and lactating animals has not been determined.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

Special precautions for use in animals:

- In the absence of available data, treatment of puppies less than 8 weeks of age and/or dogs less than 2 kg bodyweight should be based on a benefit/risk assessment by the responsible veterinarian.
- Remove only one chewable tablet at a time from the foil-backed blister card.
- Administer the chewable tablet in a manner that encourages the dog to chew rather than to swallow without chewing.
- Care should be taken to ensure that the dog consumes the complete dose; treated animals should be observed for a few minutes after administration to ensure that part of the dose is not lost or rejected.