

Warnings:

- Handle with care.
 - Keep out of reach of children, uninformed persons and animals.
 - Do not use on kittens less than 10 weeks of age.
 - Do not use in case of known hypersensitivity to the active substances or to any of the excipients.
 - Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide variety of reasons. If this is suspected, seek your veterinarian's advice and notify the registration holder.
 - Store away from food and feed.
-
- **Precautions:**
 - While occasional contact with water does not reduce the effectiveness of the collar, frequent swimming or shampooing should be avoided.
 - Wash hands with cold water after fitting the collar.
 - As with any veterinary medicinal products, do not allow small children to play with the collar or reflectors or to put them into their mouths.
 - People with sensitivity reactions to the ingredients of the collar should avoid contact with the collar.
 - Keep the bag with the collar in the outer packaging until use.
 - The collar should be worn continuously for the 8-month protection period and should be removed after the treatment period.
 - In the unlikely event of the animal eating the collar, mild gastrointestinal symptoms (e.g. loose stool) may occur.
 - Ensure that the collar is not fitted too tightly.
 - Occasionally, slight twitching may be observed in the first few days after fitting, in animals that are unused to wearing collars.
 - Slight hair loss and mild skin reactions due to the mechanical irritation of the collar may occur at the application site, but this usually resolves within 1 or 2 weeks without the need to remove the collar.
 - The safety of the product has not been established during pregnancy and lactation.